

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use DIFLUPREDNATE OPHTHALMIC EMULSION safely and effectively. See full prescribing information for DIFLUPREDNATE OPHTHALMIC EMULSION.

DIFLUPREDNATE ophthalmic emulsion, for topical ophthalmic use

Initial U.S. Approval: 2008

-----INDICATIONS AND USAGE-----

Difluprednate ophthalmic emulsion is a topical corticosteroid that is indicated for:

- The treatment of inflammation and pain associated with ocular surgery. (1.1)
- The treatment of endogenous anterior uveitis. (1.2)

-----DOSAGE AND ADMINISTRATION-----

- For the treatment of inflammation and pain associated with ocular surgery instill one drop into the conjunctival sac of the affected eye 4 times daily beginning 24 hours after surgery and continuing throughout the first 2 weeks of the postoperative period, followed by 2 times daily for a week and then a taper based on the response. (2.1)
- For the treatment of endogenous anterior uveitis instill one drop into the conjunctival sac of the affected eye 4 times daily for 14 days followed by tapering as clinically indicated. (2.2)

-----DOSAGE FORMS AND STRENGTHS-----

Difluprednate ophthalmic emulsion contains 0.05% difluprednate, as a sterile preserved ophthalmic emulsion for topical ophthalmic use only. (3)

-----CONTRAINDICATIONS-----

Difluprednate, as with other ophthalmic corticosteroids, is contraindicated in most active viral diseases of the cornea and conjunctiva including epithelial herpes simplex keratitis (dendritic keratitis), vaccinia, and varicella, and also in mycobacterial infection of the eye and fungal diseases of ocular structures. (4)

-----WARNINGS AND PRECAUTIONS-----

- Intraocular pressure (IOP) increase- Prolonged use of corticosteroids may result in glaucoma with damage to

the optic nerve, defects in visual acuity and fields of vision. If this product is used for 10 days or longer, IOP should be monitored. (5.1)

- Cataracts-Use of corticosteroids may result in posterior subcapsular cataract formation. (5.2)
- Delayed healing-The use of steroids after cataract surgery may delay healing and increase the incidence of bleb formation. In those diseases causing thinning of the cornea or sclera, perforations have been known to occur with the use of topical steroids. The initial prescription and renewal of the medication order beyond 28 days should be made by a physician only after examination of the patient with the aid of magnification such as slit lamp biomicroscopy and, where appropriate, fluorescein staining. (5.3)
- Bacterial infections-Prolonged use of corticosteroids may suppress the host response and thus increase the hazard of secondary ocular infections. In acute purulent conditions, steroids may mask infection or enhance existing infection. If signs and symptoms fail to improve after 2 days, the patient should be re-evaluated. (5.4)
- Viral infections-Employment of a corticosteroid medication in the treatment of patients with a history of herpes simplex requires great caution. Use of ocular steroids may prolong the course and may exacerbate the severity of many viral infections of the eye (including herpes simplex). (5.5)
- Fungal infections-Fungal infections of the cornea are particularly prone to develop coincidentally with long-term local steroid application. Fungus invasion must be considered in any persistent corneal ulceration where a steroid has been used or is in use. (5.6)

-----ADVERSE REACTIONS-----

To report SUSPECTED ADVERSE REACTIONS, contact Amneal Pharmaceuticals at 1-877-835-5472 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

See 17 for PATIENT COUNSELING INFORMATION.

Revised: 10/2020

FULL PRESCRIBING INFORMATION: CONTENTS*

1 INDICATIONS AND USAGE

- 1.1 Ocular Surgery
- 1.2 Endogenous Anterior Uveitis

2 DOSAGE AND ADMINISTRATION

- 2.1 Ocular Surgery
- 2.2 Endogenous Anterior Uveitis

3 DOSAGE FORMS AND STRENGTHS

4 CONTRAINDICATIONS

5 WARNINGS AND PRECAUTIONS

- 5.1 IOP Increase
- 5.2 Cataracts
- 5.3 Delayed Healing
- 5.4 Bacterial Infections
- 5.5 Viral Infections
- 5.6 Fungal Infections
- 5.7 Topical Ophthalmic Use Only
- 5.8 Contact Lens Wear

6 ADVERSE REACTIONS

- 6.1 Ocular Surgery
- 6.2 Endogenous Anterior Uveitis

8 USE IN SPECIFIC POPULATIONS

- 8.1 Pregnancy

- 8.3 Nursing Mothers
- 8.4 Pediatric Use
- 8.5 Geriatric Use

11 DESCRIPTION

12 CLINICAL PHARMACOLOGY

- 12.1 Mechanism of Action
- 12.3 Pharmacokinetics

13 NONCLINICAL TOXICOLOGY

- 13.1 Carcinogenesis, Mutagenesis, and Impairment of Fertility
- 13.2 Animal Toxicology and/or Pharmacology

14 CLINICAL STUDIES

- 14.1 Ocular Surgery
- 14.2 Endogenous Anterior Uveitis

16 HOW SUPPLIED/STORAGE AND HANDLING

17 PATIENT COUNSELING INFORMATION

- 17.1 Risk of Contamination
- 17.2 Risk of Secondary Infection
- 17.3 Contact Lens Wear

* Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION

1 INDICATIONS AND USAGE

1.1 Ocular Surgery

Difluprednate ophthalmic emulsion, 0.05%, a topical corticosteroid, is indicated for the treatment of inflammation and pain associated with ocular surgery.

1.2 Endogenous Anterior Uveitis

Difluprednate ophthalmic emulsion, 0.05% is also indicated for the treatment of endogenous anterior uveitis.

2 DOSAGE AND ADMINISTRATION

2.1 Ocular Surgery

Instill one drop into the conjunctival sac of the affected eye 4 times daily beginning 24 hours after surgery and continuing throughout the first 2 weeks of the postoperative period, followed by 2 times daily for a week and then a taper based on the response.

2.2 Endogenous Anterior Uveitis

Instill one drop into the conjunctival sac of the affected eye 4 times daily for 14 days followed by tapering as clinically indicated.

3 DOSAGE FORMS AND STRENGTHS

Difluprednate ophthalmic emulsion contains 0.05% difluprednate as a sterile preserved emulsion for topical ophthalmic administration.

4 CONTRAINDICATIONS

The use of difluprednate, as with other ophthalmic corticosteroids, is contraindicated in most active viral diseases of the cornea and conjunctiva including epithelial herpes simplex keratitis (dendritic keratitis), vaccinia, and varicella, and also in mycobacterial infection of the eye and fungal disease of ocular structures.

5 WARNINGS AND PRECAUTIONS

5.1 IOP Increase

Prolonged use of corticosteroids may result in glaucoma with damage to the optic nerve, defects in visual acuity and fields of vision. Steroids should be used with caution in the presence of glaucoma. If this product is used for 10 days or longer, intraocular pressure should be monitored.

5.2 Cataracts

Use of corticosteroids may result in posterior subcapsular cataract formation.

5.3 Delayed Healing

The use of steroids after cataract surgery may delay healing and increase the incidence of bleb formation. In those diseases causing thinning of the cornea or sclera, perforations have been known to occur with the use of topical steroids. The initial prescription and renewal of the medication order beyond 28 days should be made by a physician only after examination of the patient with the aid of magnification such as slit lamp biomicroscopy and, where appropriate, fluorescein staining.

5.4 Bacterial Infections

Prolonged use of corticosteroids may suppress the host response and thus increase the hazard of secondary ocular infections. In acute purulent conditions, steroids may mask infection or enhance existing infection. If signs and symptoms fail to improve after 2 days, the patient should be re-evaluated.

5.5 Viral Infections

Employment of a corticosteroid medication in the treatment of patients with a history of herpes simplex requires great caution. Use of ocular steroids may prolong the course and may exacerbate the severity of many viral infections of the eye (including herpes simplex).

5.6 Fungal Infections

Fungal infections of the cornea are particularly prone to develop coincidentally with long-term local steroid application. Fungus invasion must be considered in any persistent corneal ulceration where a steroid has been used or is in use. Fungal culture should be taken when appropriate.

5.7 Topical Ophthalmic Use Only

Difluprednate is not indicated for intraocular administration.

5.8 Contact Lens Wear

Difluprednate ophthalmic emulsion should not be instilled while wearing contact lenses. Remove contact lenses prior to instillation of difluprednate ophthalmic emulsion. The preservative in difluprednate ophthalmic emulsion may be absorbed by soft contact lenses. Lenses may be reinserted after 10 minutes following administration of difluprednate ophthalmic emulsion.

6 ADVERSE REACTIONS

The following serious reactions are found elsewhere in the labeling:

- Elevated intraocular pressure [*see Warnings and Precautions (5.1)*]
- Posterior subcapsular cataract formation [*see Warnings and Precautions (5.2)*]
- Secondary ocular infection [*see Warnings and Precautions (5.4)*]
- Perforation of the globe [*see Warnings and Precautions (5.3)*]

6.1 Ocular Surgery

Ocular adverse reactions occurring in 5% to 15% of subjects in clinical studies with difluprednate included corneal edema, ciliary and conjunctival hyperemia, eye pain, photophobia, posterior capsule opacification, anterior chamber cells, anterior chamber flare, conjunctival edema, and blepharitis. Other ocular adverse reactions occurring in 1% to 5% of subjects included reduced visual acuity, punctate keratitis, eye

inflammation, and iritis. Ocular adverse reactions occurring in < 1% of subjects included application site discomfort or irritation, corneal pigmentation and striae, episcleritis, eye pruritis, eyelid irritation and crusting, foreign body sensation, increased lacrimation, macular edema, sclera hyperemia, and uveitis. Most of these reactions may have been the consequence of the surgical procedure.

6.2 Endogenous Anterior Uveitis

A total of 200 subjects participated in the clinical trials for endogenous anterior uveitis, of which 106 were exposed to difluprednate. The most common adverse reactions of those exposed to difluprednate occurring in 5% to 10% of subjects included blurred vision, eye irritation, eye pain, headache, increased IOP, iritis, limbal and conjunctival hyperemia, punctate keratitis, and uveitis. Adverse reactions occurring in 2% to 5% of subjects included anterior chamber flare, corneal edema, dry eye, iridocyclitis, photophobia, and reduced visual acuity.

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Teratogenic Effects

Pregnancy Category C

Difluprednate has been shown to be embryotoxic (decrease in embryonic body weight and a delay in embryonic ossification) and teratogenic (cleft palate and skeletal anomalies) when administered subcutaneously to rabbits during organogenesis at a dose of 1 to 10 mcg/kg/day. The no-observed-effect-level (NOEL) for these effects was 1 mcg/kg/day, and 10 mcg/kg/day was considered to be a teratogenic dose that was concurrently found in the toxic dose range for fetuses and pregnant females. Treatment of rats with 10 mcg/kg/day subcutaneously during organogenesis did not result in any reproductive toxicity, nor was it maternally toxic. At 100 mcg/kg/day after subcutaneous administration in rats, there was a decrease in fetal weights and delay in ossification, and effects on weight gain in the pregnant females. It is difficult to extrapolate these doses of difluprednate to maximum daily human doses of difluprednate, since difluprednate is administered topically with minimal systemic absorption, and difluprednate blood levels were not measured in the reproductive animal studies. However, since use of difluprednate during human pregnancy has not been evaluated and cannot rule out the possibility of harm, difluprednate should be used during pregnancy only if the potential benefit justifies the potential risk to the embryo or fetus.

8.3 Nursing Mothers

It is not known whether topical ophthalmic administration of corticosteroids could result in sufficient systemic absorption to produce detectable quantities in breast milk. Systemically administered corticosteroids appear in human milk and could suppress growth, interfere with endogenous corticosteroid production, or cause other untoward effects. Caution should be exercised when difluprednate is administered to a nursing woman.

8.4 Pediatric Use


Difluprednate was evaluated in a 3-month, multicenter, double-masked trial in 79 pediatric patients (39 difluprednate; 40 prednisolone acetate) 0 to 3 years of age for the treatment of inflammation following cataract surgery. A similar safety profile was observed in pediatric patients comparing difluprednate to prednisolone acetate ophthalmic suspension, 1%.

8.5 Geriatric Use

No overall differences in safety or effectiveness have been observed between elderly and younger patients.

11 DESCRIPTION

Difluprednate ophthalmic emulsion, 0.05% is a sterile, topical anti-inflammatory corticosteroid for ophthalmic use. The chemical name is (6 α ,11 β)-21-(Acetyloxy)-6,9-difluoro-11-hydroxy-17-(1-oxobutoxy)pregna-1,4-diene-3,20-dione. Difluprednate is represented by the following structural formula:



Difluprednate has a molecular weight of 508.55, and the molecular formula is C₂₇H₃₄F₂O₇.

Difluprednate (micronized) is a white to creamy-white crystalline powder. It is freely soluble in acetonitrile and in chloroform, soluble in methanol, in ethanol and in dioxane; slightly soluble in ether; practically insoluble in water.

Each mL contains:

Active: difluprednate 0.5 mg (0.05%)

Inactive: boric acid, castor oil, glycerin, polysorbate 80, water for injection, sodium acetate, edetate disodium, sodium hydroxide (to adjust the pH to 5.2 to 5.8). The emulsion is essentially isotonic with a tonicity of 304 to 411 mOsmol/kg

Preservative: sorbic acid 0.1%

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

Corticosteroids inhibit the inflammatory response to a variety of inciting agents and may delay or slow healing. They inhibit edema, fibrin deposition, capillary dilation, leukocyte migration, capillary proliferation, fibroblast proliferation, deposition of collagen, and scar formation associated with inflammation. There is no generally accepted explanation for the mechanism of action of ocular corticosteroids. However, corticosteroids are thought to act by the induction of phospholipase A2 inhibitory proteins, collectively called lipocortins. It is postulated that these proteins control the biosynthesis of potent mediators of inflammation such as prostaglandins and leukotrienes by inhibiting the release of their common precursor arachidonic acid. Arachidonic acid is released from membrane phospholipids by phospholipase A2.

Difluprednate is structurally similar to other corticosteroids.

12.3 Pharmacokinetics

Diffuprednate undergoes deacetylation *in vivo* to 6 α , 9-difluoroprednisolone 17-butyrate (DFB), an active metabolite of difluprednate.

Clinical pharmacokinetic studies of difluprednate after repeat ocular instillation of 2 drops of difluprednate (0.01% or 0.05%) four times per day for 7 days showed that DFB levels in blood were below the quantification limit (50 ng/mL) at all time points for all subjects, indicating the systemic absorption of difluprednate after ocular instillation of difluprednate is limited.

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, and Impairment of Fertility

Difluprednate was not genotoxic *in vitro* in the Ames test, and in cultured mammalian cells CHL/IU (a fibroblastic cell line derived from the lungs of newborn female Chinese hamsters). An *in vivo* micronucleus test of difluprednate in mice was also negative. Treatment of male and female rats with subcutaneous difluprednate up to 10 mcg/kg/day prior to and during mating did not impair fertility in either gender. Long term studies have not been conducted to evaluate the carcinogenic potential of difluprednate.



13.2 Animal Toxicology and/or Pharmacology

In multiple studies performed in rodents and non-rodents, subchronic and chronic toxicity tests of difluprednate showed systemic effects such as suppression of body weight gain; a decrease in lymphocyte count; atrophy of the lymphatic glands and adrenal gland; and for local effects, thinning of the skin; all of which were due to the pharmacologic action of the molecule and are well known glucocorticosteroid effects. Most, if not all of these effects were reversible after drug withdrawal. The NOEL for the subchronic and chronic toxicity tests were consistent between species and ranged from 1 to 1.25 mcg/kg/day.

14 CLINICAL STUDIES

14.1 Ocular Surgery

Clinical efficacy was evaluated in 2 randomized, double-masked, placebo-controlled trials in which subjects with an anterior chamber cell grade \geq "2" (a cell count of 11 or higher) after cataract surgery were assigned to difluprednate or placebo (vehicle) following surgery. One drop of difluprednate ophthalmic emulsion or vehicle was self instilled either 2 times per day or 4 times per day for 14 days, beginning the day after surgery. The presence of complete clearing (a cell count of 0) was assessed 3, 8 and 15 days post-surgery using a slit lamp binocular microscope. In the intent-to-treat analyses of both studies, a significant benefit was seen in the 4 times per day (QID) difluprednate-treated group in ocular inflammation, at days 8 and 15, and reduction of pain, at days 3, 8 and 15, when compared with placebo. The consolidated clinical trial results are provided below.



14.2 Endogenous Anterior Uveitis

Clinical efficacy was evaluated in two randomized, double masked active controlled trials in which patients who presented with endogenous anterior uveitis were treated with either difluprednate 4 times daily or

prednisolone acetate ophthalmic suspension, 1%, 8 times daily for 14 days. Both studies demonstrated that difluprednate was equally effective as prednisolone acetate ophthalmic suspension, 1% in treating subjects with endogenous anterior uveitis.

Mean Change from Baseline in Anterior Chamber Cell Grade*

Study 1 time point	Difluprednate N=57	Prednisolone Acetate N=53	Difference [†] (95% CI)
Baseline	2.6	2.5	0.0 (-0.22, 0.28)
Day 3	-1.0	-1.0	-0.1 (-0.35, 0.25)
Day 7	-1.6	-1.5	-0.0 (-0.31, 0.25)
Day 14	-2.0	-1.8	-0.2 (-0.46, 0.10)
Day 21	-2.2	-1.9	-0.3 (-0.53, 0.01)
Day 28	-2.2	-2.1	-0.1 (-0.37, 0.18)
Day 35	-2.1	-2.0	-0.1 (-0.39, 0.20)
Day 42	-2.1	-2.1	0.0 (-0.27, 0.34)
Study 2 time point	Difluprednate N=50	Prednisolone Acetate N=40	Difference [†] (95% CI)
Baseline	2.4	2.4	0.0 (-0.21, 0.29)
Day 3	-0.9	-0.9	-0.0 (-0.34, 0.25)
Day 7	-1.7	-1.6	-0.1 (-0.35, 0.21)
Day 14	-1.9	-1.8	-0.1 (-0.34, 0.20)
Day 21	-2.0	-2.0	0.0 (-0.25, 0.28)
Day 28	-2.0	-2.0	0.0 (-0.21, 0.26)
Day 35	-2.1	-2.0	-0.1 (-0.32, 0.16)
Day 42	-2.0	-1.9	-0.1 (-0.36, 0.24)

* with 5 grades: 0 = 0 cells; 1 = 1 to 10 cells; 2 = 11 to 20 cells; 3 = 21 to 50 cells; and 4 = > 50 cells.
[†] adjusted for baseline AC cell grade and study center and based on ITT dataset with LOCF for missing data.

16 HOW SUPPLIED/STORAGE AND HANDLING

Difluprednate Ophthalmic Emulsion, **0.05%** is a sterile, white milky aqueous topical ophthalmic emulsion and supplied in white LDPE opaque bottle with a LDPE natural nozzle and HDPE pink cap.

It is available as follows:

5 mL in 5 mL Container: NDC 69238-1380-3

Storage and Handling

Store at 15° to 25°C (59° to 77°F). Do not freeze. Protect from light. When not in use, keep the bottles in the protective carton.

17 PATIENT COUNSELING INFORMATION

17.1 Risk of Contamination

This product is sterile when packaged. Advise patients not to allow the dropper tip to touch any surface, as this may contaminate the emulsion.

Use of the same bottle for both eyes is not recommended with topical eye drops that are used in association with surgery.

17.2 Risk of Secondary Infection

If pain develops, or if redness, itching, or inflammation becomes aggravated, advise patients to consult a physician.

17.3 Contact Lens Wear

Difluprednate ophthalmic emulsion should not be instilled while wearing contact lenses. Advise patients to remove contact lenses prior to instillation of difluprednate ophthalmic emulsion. The preservative in difluprednate ophthalmic emulsion may be absorbed by soft contact lenses. Lenses may be reinserted after 10 minutes following administration of difluprednate ophthalmic emulsion.

Manufactured by:

Amneal Pharmaceuticals Pvt. Ltd.,

Parenteral Unit

Ahmedabad 382213, INDIA

Distributed by:

Amneal Pharmaceuticals LLC

Bridgewater, NJ 08807

Rev. 10-2020-01